



Medical Management of Chronic Pancreatitis in Children

Head of Pediatric Gastroenterology and Hepatology Research Center

Pejman Rohani MD

Associate Professor

ESPGHAN Member

PIBD SIG Group

Pediatric Center of Excellence, Children's Medical Center

TUMS



CP requires imaging findings characteristic of and consistent with CP (specifically, radiographically evident calcifications, and pancreatic duct irregularities, such as strictures and dilations) along with 1 of the following 3: abdominal pain consistent with pancreatic origin, evidence of exocrine pancreatic insufficiency (EPI), or evidence of endocrine pancreatic insufficiency, or a pancreatic biopsy specimen demonstrating histological evidence of CP

In recent years, studies have shown a relatively high rate of associated genetic mutations, with positive genetic testing in 64% to 70% of pediatric chronic pancreatitis cases.

Chronic ductal obstruction from anatomic abnormalities, medication induced pancreatitis, autoimmune pancreatitis, and inborn errors of metabolism should also be considered.

Currently, the known common mutations include the cationic trypsinogen gene, *PRSS1*, along with the cystic fibrosis transmembrane conductance regulator (*CFTR*) and serine peptidase inhibitor Kazal type 1 (*SPINK1*) mutations.⁸⁷ Other associated mutations include chymotrypsin C (caldecrin) gene (*CTRC*) and the calcium-sensing receptor gene (*CASR*), among other associated genetic mutations.

Etiology of Chronic Pancreatitis

Toxic-metabolic

Medications (see Box 82.1)

Hypercalcemia

Hyperlipidemia

Post-graft-versus-host disease

Toxins

Organic compounds

Genetic

Autosomal dominant

PRSS1 mutations

Autosomal recessive/modifiers

CFTR mutations

SPINK1 mutations

CPA1 mutations

CTRC mutations

Autoimmune

Isolated autoimmune pancreatitis

Syndromic autoimmune pancreatitis

Inflammatory bowel disease associated

Anatomic

See Box 82.1

Severe acute pancreatitis

Postnecrotic

Vascular disease/ischemic

Postirradiation

Indeterminate

Imaging

Dietary Considerations for Children With Chronic Pancreatitis

**Use of Pancreatic Enzyme Replacement Therapy for
Exocrine Pancreatic Insufficiency in Children With Chronic
Pancreatitis**

**Pancreatogenic Diabetes Mellitus in Children With
Chronic Pancreatitis**

Medical Management of Pain in Chronic Pancreatitis

Sequelae of Chronic Pancreatitis

Imaging

In general terms, the roles of imaging in CP are to:

- Contribute to/establish the initial diagnosis of CP;
- Stage and monitor disease, including complications;
- Assess for superimposed AP;
- Identify potential etiologies of CP;
- Identify findings that might herald endocrine or exocrine dysfunction;
- Characterize secretory (exocrine) function;
- Plan for intervention.

Although findings of CP may be identified on ultrasound or CT, MRI/magnetic resonance cholangiopancreatography (MRI/MRCP) is favored for the diagnosis and characterization of CP given its superiority in visualizing parenchymal and duct changes

Ultrasound examination

Ultrasound examinations are ideally performed fasting (4 hours) to reduce bowel gas that can obscure the pancreas and to distend the gallbladder. When the patient is able, drinking water immediately prior to the examination to distend the stomach with fluid and displace gastric air may provide an improved acoustic

window for imaging the pancreas.

A complete ultrasound examination should assess pancreatic size, contour, echogenicity, pancreatic duct diameter (and for duct filling defects), and should assess for peripancreatic edema and pancreatic or peripancreatic fluid collections. In addition, the gallbladder should be assessed for calculi (an etiology of pancreatitis), and the biliary tree should be assessed for dilation and calculi.

Color Doppler along with gray-scale imaging can evaluate the peripancreatic vascular structures for complications, such as splenic vein or portal vein thrombosis and can assess vascularity of the pancreas.

Computed Tomography

Use of IV contrast material is recommended when performing CT for CP.

IV contrast material allows optimal assessment of the pancreatic parenchyma and allows evaluation of the peripancreatic vessels for patency.

For adults, multiphase protocols that include an unenhanced phase, parenchymal/arterial phase, and portal venous phase have been recommended.

No such recommendations exist for pediatrics but given the relative infrequency of calcifications in pediatric pancreatitis, an unenhanced phase is likely unnecessary.

As with AP, a parenchymal/arterial phase can be useful if clinical questions relate to the arteries but a single portal venous phase examination is generally sufficient to characterize CP in children.

CT with IV contrast material provides excellent assessment of the pancreatic parenchyma, allowing identification of features of ARP and CP, particularly calcifications and pancreatic atrophy.

CT can also identify congenital anomalies, such as annular pancreas and can assess for superimposed acute pancreatitis and complications of pancreatitis, including established vascular collaterals because of chronic/established thrombosis.

CT (and MRI) outperform ultrasound to define vascular anatomy relevant to surgical planning. CT is, however, limited by suboptimal visualization of the pancreatic and biliary ducts.

Magnetic Resonance Imaging/Magnetic Resonance Cholangiopancreatography

MRI and MRCP have the benefits of providing information on both parenchymal and duct changes of CP but are limited in their ability to visualize calcifications.

Adult and pediatric data suggest that the sensitivity of MRCP to detect pancreatic duct abnormalities may be improved by the administration of secretin.

Theoretically secretin distends the pancreatic duct and may allow for earlier detection of side branch-ectasias and provide information on exocrine function by quantifying duodenal filling

Chronic Pancreatitis Summary Statements and Recommendations

Initial Diagnosis

MRI is the recommended modality for imaging of suspected CP.

a. This recommendation reflects inadequate characterization of findings (particularly duct findings) of CP by transabdominal ultrasound. This also reflects the superior soft tissue contrast of MRI, which allows characterization of both parenchyma and duct findings of CP.

Assessment for Superimposed Acute Pancreatitis

When imaging is needed to assess a suspected or known episode of AP in a child with CP, transabdominal ultrasound is the preferred first-line imaging modality.

a. This recommendation reflects the availability and portability of ultrasound and the role of ultrasound in identifying biliary causes of AP.

b. Note: A negative ultrasound does not exclude AP (low to moderate sensitivity).

If ultrasound is negative for AP in a child with CP and an imaging diagnosis of AP is needed, either CT or

Intervention Planning

CT or MRI are recommended for planning of endoscopic or

surgical interventions in a patient with known CP.

a. This recommendation is based on the large field of view afforded by CT and MRI, optimal characterization of the degree of organization of fluid collections, and optimal characterization of the peripancreatic vasculature.


b. Note: When the intervention will target the duct, MRI is favored over CT.

Serial Monitoring

MRI is recommended for clinically indicated serial imaging of CP.


a. This recommendation reflects the optimal soft tissue contrast of MRI, no need for intravenous contrast material (in most cases), and the lack of associated ionizing radiation.

b. Note: In the child who requires sedation for MRI, risks of sedation must be balanced with the need for serial imaging.



Proposed analgesic ladder for the treatment of pain in children with chronic pancreatitis

EVALUATING EXOCRINE PANCREATIC INSUFFICIENCY (EPI) AND INITIATING PANCREATIC ENZYME REPLACEMENT THERAPY (PERT)



This clinical pathway is based upon publicly available medical evidence and/or a consensus of medical practitioners at the time of publication. It is intended to be a guide for general practitioners and may need to be adapted for each specific patient based on the practitioner's professional judgment, consideration of any unique circumstances, the needs of each patient and their family, and/or the availability of various resources at the health care institution where the patient is located.

Conditions with High Risk of EPI

Acquired Pancreatic Insufficiency

- Chronic Pancreatitis
- Status-post Pancreatectomy
- Chemotherapy
- Trauma
- Autoimmune/inflammatory disorders

Symptoms Suggestive of EPI

- Poor weight gain/linear growth
- Steatorrhea
- Loose/frequent stools after meals/snacks
- Foul-smelling stools
- Flatulence
- Abdominal pain associated with eating
- Voracious Appetite
- Rectal Prolapse

Elastase:

- Requires a single spot sample
- Results may be inaccurate if stool is watery
- Differences between monoclonal and polyclonal testing

Coefficient of Fat Absorption:

- Requires 72-hour high-fat diet and stool collection

*Universal agreement at the ends of the reference range; different reference ranges used by different labs. Not a continuous scale